




# Increase patient care. Decrease your costs.

The leader in the fight against premature birth, March of Dimes, states that:




**\$26 billion**

is spent caring for babies in the NICU.



**93%** of premature infants are diagnosed with **respiratory distress syndrome (RDS)** and require ventilator support.

Use the *one* high frequency ventilator designed for *all* of your neonatal patients!



- + Lower hospital cost<sup>1,2</sup>
- + As little as two days on HFOV before extubation<sup>3,7</sup>
- + Decreased duration of oxygen therapy<sup>3,6</sup>
- + Decreased nCPAP duration<sup>2,6</sup>
- + Decreased chronic lung disease (CLD)<sup>1-5</sup>
- + Discharge up to 24 days earlier<sup>3</sup>



"For the subgroup of 11 trials involving 1,737 infants that used high frequency piston oscillators, there was a significant reduction in CLD in the HFOV group."

The Cochrane Collaboration<sup>®</sup> 2015

Can you afford *not* to use the 3100A HFOV?

For educational resources, including webinars about lung recruitment, visit [vyaire.com](http://vyaire.com)



#### References

1 Gerstmann D, Wood K, Miller A, et al. Childhood outcome after early high-frequency oscillatory ventilation for neonatal respiratory distress syndrome. *Pediatrics*. 2001;108(3):617-23. 2 Courtney S, Durand D, Asselin J, et al. High-Frequency Oscillatory Ventilation versus Conventional Mechanical Ventilation for Very-Low-Birth-Weight Infants. *NEJM*. 2002;347:643-652. 3 Salvo V, Zimmermann L, Gavilanes A, et al. First intention high-frequency oscillatory and conventional mechanical ventilation in premature infants without antenatal glucocorticoid prophylaxis. *Pediatric Critical Care Med*. 2012;13:72-79. 4 Clark R, Gerstmann D, Null D, et al. Prospective randomized comparison of high frequency oscillatory and conventional ventilation in respiratory distress syndrome. *Pediatrics*. 1992;89:5-12. 5 Gerstmann D, Minton S, Stoddard R, et al. The provo multicenter early high frequency oscillatory ventilation trial: improved pulmonary and clinical outcome in respiratory distress syndrome. *Pediatrics*. 1996;98:1044-57. 6 Rimensberger P, Beghetti M, Hanquinet S, et al. First Intention High-Frequency Oscillation with Early Lung Volume Optimization Improves Pulmonary Outcome in Very Low Birth Weight Infants with Respiratory Distress Syndrome. *Pediatrics*. 2000;105:6. 7 De Jaegere A, van Veenendaal M, Michiels A, van Kaam A. Lung recruitment using oxygenation during open lung high-frequency ventilation in preterm infants. *Respir Care Clin N Am*, 2006. 174:639645.

\* Based on U.S. statistics from marchofdimes.org

Vyaire Medical  
Mettawa, IL  
[vyaire.com](http://vyaire.com)

